

## Two Theorems on Finite Hilbert Transforms\*

MARVIN ROSENBLUM AND JAMES ROVNYAK

*University of Virginia, Charlottesville, Virginia 22903**Submitted by Peter D. Lax*

Our results are motivated by a theorem of C. Loewner. Let  $f(x)$  be a real valued function which is continuously differentiable on an interval  $(a, b)$ . The function  $[f(x) - f(t)]/(x - t)$ , extended by continuity to the square  $(a, b) \times (a, b)$ , is called positive definite if

$$\sum_{j,k=1}^n \frac{f(x_k) - f(x_j)}{x_k - x_j} c_j c_k^* \geq 0 \quad (1)$$

for all finite sets of points  $x_1, \dots, x_n$  in  $(a, b)$  and complex numbers  $c_1, \dots, c_n$ . Loewner [7] showed that  $[f(x) - f(t)]/(x - t)$  is positive definite if and only if  $f(x)$  is the restriction to  $(a, b)$  of a function  $f(z)$  which is analytic in the upper and lower half-planes and across  $(a, b)$  and satisfies  $\operatorname{Im} f(z) \geq 0$  for  $y > 0$ . For other treatments and related results see [1, 5, 6, 8, 12]. The first of our two theorems is an extension of Loewner's theorem. The second, a dual result, includes as a special case a characterization of nonnegative integrable functions  $g(x)$  on  $(a, b)$  such that

$$\int_a^b \left| PV \frac{1}{\pi} \int_a^b \frac{\gamma(t)}{t - x} dt \right|^2 g(x) dx \leq \int_a^b |\gamma(x)|^2 g(x) dx$$

for any sufficiently large class of functions  $\gamma(x)$ .

It is sufficient to treat the case of a symmetric interval  $(-a, a)$ . By a *locally finite signed measure* (or "*lfs measure*") on  $(-a, a)$  we mean a real valued set function which is finite and completely additive on the Borel subsets of  $(-a, a)$  that are at a positive distance from the end points. The complex conjugate of a number  $c$  or function  $\gamma$  is written  $c^*$  or  $\gamma^*$ , respectively.

For any  $\gamma \in L^1(-\infty, \infty)$  the conjugate function  $\tilde{\gamma}$  is defined a.e. on  $(-\infty, \infty)$  by

$$\tilde{\gamma}(x) = PV \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\gamma(t)}{t - x} dt,$$

\* Research supported by NSF Grant No. GP-31483X.

where  $PV$  indicates that the integral is taken as a Cauchy principal value. Given  $\gamma \in L^1(-a, a)$  we set  $\gamma = 0$  off  $(-a, a)$  and regard  $\tilde{\gamma}$  as defined a.e. on  $(-\infty, \infty)$ , though we are usually interested only in its restriction to  $(-a, a)$ . Let  $\phi \in L^p(-a, a)$ ,  $\psi \in L^q(-a, a)$  where  $p > 1$ ,  $q > 1$ . Then  $\tilde{\phi} \in L^p(-a, a)$ ,  $\tilde{\psi} \in L^q(-a, a)$ . If  $p^{-1} + q^{-1} \leq 1$  then

$$\int_{-a}^a (\phi \tilde{\psi} + \tilde{\phi} \psi) dt = 0. \quad (2)$$

If  $p^{-1} + q^{-1} < 1$  then [10, p. 169]

$$(\phi \tilde{\psi} + \tilde{\phi} \psi) = \tilde{\phi} \tilde{\psi} - \phi \psi \quad (3)$$

a.e. on  $(-a, a)$  (and indeed a.e. on  $(-\infty, \infty)$ ).

Let  $\mathcal{D}_a$  be the class of continuous functions  $\gamma$  on  $(-\infty, \infty)$  which vanish off some closed subinterval of  $(-a, a)$  such that  $\tilde{\gamma}$  is everywhere defined and continuous on  $(-\infty, \infty)$ .

**PROBLEM 1.** Characterize the class  $\mathcal{U}$  of lfs measures  $\mu$  on  $(-a, a)$  such that

$$\operatorname{Re} \int_{(-a, a)} \tilde{\gamma} \gamma^* d\mu \leq 0 \quad (4)$$

for all  $\gamma \in \mathcal{D}_a$ .

We first connect this problem with Loewner's theorem.

**PROPOSITION 1.** Let  $d\mu = f dt$  where  $f$  is a real valued function which is integrable over every closed subinterval of  $(-a, a)$ . Then  $\mu \in \mathcal{U}$  if and only if

$$\lim_{\epsilon \downarrow 0} \int_{-a}^a \int_{-a}^a \frac{f(x) - f(t)}{x - t} \delta_\epsilon(t - x) \gamma(t) \gamma^*(x) dt dx \geq 0 \quad (5)$$

for all  $\gamma \in \mathcal{D}_a$ , where  $\delta_\epsilon(t) = 1$  or  $0$  according as  $|t| > \epsilon$  or  $|t| < \epsilon$ . The limit exists for all  $\gamma \in \mathcal{D}_a$  whether or not  $\mu \in \mathcal{U}$ .

If  $f(x)$  is continuously differentiable on  $(-a, a)$ , then (5) is equivalent to (1). Therefore any characterization of the class  $\mathcal{U}$  must include Loewner's theorem.

**LEMMA 1.** Let  $\gamma$  be continuous on  $(-\infty, \infty)$  and vanish off some closed subinterval of  $(-a, a)$ . If  $\tilde{\gamma}$  is equal a.e. to a function which is continuous on  $(-\infty, \infty)$  then  $\gamma \in \mathcal{D}_a$ . In fact, under these conditions

$$\tilde{\gamma}(x) = \lim_{\epsilon \downarrow 0} \frac{1}{\pi} \int_{-a}^a \frac{\gamma(t)}{t - x} \delta_\epsilon(t - x) dt$$

uniformly on  $(-\infty, \infty)$ .

*Proof.* Let  $\tilde{\gamma} = \beta$  a.e. where  $\beta$  is continuous on  $(-\infty, \infty)$ . Following Titchmarsh [9, pp. 122–125] we define

$$V(x, y) = -\frac{1}{\pi} \int_{-a}^a \frac{t-x}{(t-x)^2 + y^2} \gamma(t) dt$$

and argue that

$$V(x, y) = -\frac{y}{\pi} \int_{-\infty}^{\infty} \frac{\beta(t)}{(t-x)^2 + y^2} dt$$

for all real  $x$  and  $y > 0$ . Since  $\beta$  is uniformly continuous on  $(-\infty, \infty)$ ,

$$\beta(x) = -\lim_{y \downarrow 0} V(x, y)$$

uniformly on  $(-\infty, \infty)$ . Next write

$$\begin{aligned} V(x, y) + \frac{1}{\pi} \int_{-a}^a \frac{\gamma(t)}{t-x} \delta_y(t-x) dt \\ &= V(x, y) + \frac{1}{\pi} \int_y^{\infty} \frac{\gamma(x+t) - \gamma(x-t)}{t} dt \\ &= -\frac{1}{\pi} \int_0^y \frac{t}{t^2 + y^2} [\gamma(x+t) - \gamma(x-t)] dt \\ &\quad + \frac{y^2}{\pi} \int_y^{\infty} \frac{\gamma(x+t) - \gamma(x-t)}{(t^2 + y^2)t} dt. \end{aligned}$$

Let  $\epsilon > 0$  be given. Let  $|\gamma(t)| \leq M$  for all real  $t$ , and choose  $y_0 > 0$  by uniform continuity so that  $|\gamma(x+t) - \gamma(x-t)| < \epsilon$  whenever  $0 \leq t \leq y_0$ . Choose  $y_1$  so that  $0 < y_1 \leq y_0$  and  $My_1^2/(\pi y_0^2) < 2\epsilon/3$ . Then if only  $0 < y < y_1$ ,

$$\begin{aligned} &\left| V(x, y) + \frac{1}{\pi} \int_{-a}^a \frac{\gamma(t)}{t-x} \delta_y(t-x) dt \right| \\ &\leq \frac{\epsilon}{2\pi y} \int_0^y dt + \frac{\epsilon y^2}{\pi} \int_y^{y_0} \frac{dt}{(t^2 + y^2)t} + \frac{2My^2}{\pi} \int_{y_0}^{\infty} \frac{dt}{t^3} \\ &< \epsilon \left( \frac{1}{2\pi} + \frac{y^2}{\pi} \int_y^{\infty} \frac{dt}{t^3} \right) + My^2/(\pi y_0^2) \\ &< \epsilon \end{aligned}$$

uniformly in  $x$ .

*Proof of Proposition 1.* For any  $\gamma \in \mathcal{D}_a$ ,

$$\begin{aligned} 2 \operatorname{Re} \int_{(-a, a)} \tilde{\gamma} \gamma^* d\mu &= \int_{-a}^a \tilde{\gamma}(x) \gamma^*(x) f(x) dx + \int_{-a}^a \tilde{\gamma}^*(t) \gamma(t) f(t) dt \\ &= \lim_{\epsilon \downarrow 0} \left[ \int_{-a}^a \frac{1}{\pi} \int_{-a}^a \frac{\gamma(t)}{t-x} \delta_\epsilon(t-x) dt \gamma^*(x) f(x) dx \right. \\ &\quad \left. + \int_{-a}^a \frac{1}{\pi} \int_{-a}^a \frac{\gamma^*(x)}{x-t} \delta_\epsilon(x-t) dx \gamma(t) f(t) dt \right] \\ &= -\lim_{\epsilon \downarrow 0} \frac{1}{\pi} \int_{-a}^a \int_{-a}^a \frac{f(x) - f(t)}{x-t} \delta_\epsilon(t-x) \gamma(t) \gamma^*(x) dt dx. \end{aligned}$$

The change of order of integration and passage to the limit is justified by Lemma 1. The result follows.

**THEOREM 1.** *An lfs measure  $\mu$  on  $(-a, a)$  belongs to the class  $\mathcal{U}$  if and only if  $d\mu = f dx$ , where*

$$f(z) = p + \int_{-1/a}^{1/a} \frac{z}{1-zt} d\alpha(t), \quad (6)$$

$z \notin (-\infty, -a] \cup [a, \infty)$ , for some real constant  $p$  and nondecreasing function  $\alpha(t)$  on  $[-1/a, 1/a]$ .

*Proof of necessity.* To begin suppose  $a > 1$ . The Chebychev polynomials  $T_j(x)$ ,  $U_j(x)$  can be defined by

$$T_j(\cos \theta) = \cos j\theta, \quad U_j(\cos \theta) = \sin(j+1)\theta/\sin \theta$$

for all  $j \geq 0$ . A standard formula [3, p. 254] gives the conjugate pair

$$\begin{aligned} \gamma_j(x) &= \begin{cases} U_j(x)(1-x^2)^{1/2}, & |x| \leq 1 \\ 0, & |x| > 1 \end{cases} \\ \tilde{\gamma}_j(x) &= \begin{cases} -T_{j+1}(x), & |x| \leq 1 \\ \text{---}, & |x| > 1 \end{cases} \end{aligned}$$

for each  $j \geq 0$ . Also  $\gamma_j \in \mathcal{D}_a$ ,  $j \geq 0$ . For  $j = 0$  this is proved by computing  $\tilde{\gamma}_0$  on all of  $(-\infty, \infty)$  (see [3, p. 246]) and noting that it is continuous. The general case follows from the fact that  $\mathcal{D}_a$  is closed under multiplication by  $x$ .

Now substitute

$$\begin{cases} \gamma(x) = \sum_{j=0}^n c_j U_j(x) (1-x^2)^{1/2} \chi_{(-1,1)}(x) \\ \tilde{\gamma}(x) \chi_{(-1,1)}(x) = -\sum_{j=0}^n c_j T_{j+1}(x) \chi_{(-1,1)}(x) \end{cases} \quad (7)$$

into (4) to get

$$\sum_{j,k=0}^n \int_{(-1,1)} [U_k(t) T_{j+1}(t) + U_j(t) T_{k+1}(t)] (1-t^2)^{1/2} d\mu(t) c_j c_k^* \geq 0$$

and hence

$$\sum_{j,k=0}^n \int_{(-1,1)} U_{j+k+1}(t) (1-t^2)^{1/2} d\mu(t) c_j c_k^* \geq 0$$

for any complex numbers  $c_0, \dots, c_n$ . By the theory of the Hamburger moment problem [11, pp. 129–136] there exists a nondecreasing function  $\alpha_0(t)$  such that

$$\int_{(-1,1)} U_{j+1}(t) (1-t^2)^{1/2} d\mu(t) = \int_{-\infty}^{\infty} t^j d\alpha_0(t)$$

for all  $j \geq 0$ . Since

$$\begin{aligned} \int_{-\infty}^{\infty} t^{2j} d\alpha_0(t) &\leq \int_{(-1,1)} |U_{2j+1}(t)| (1-t^2)^{1/2} d\mu(t) \\ &\leq (2j+2) \int_{(-1,1)} (1-t^2)^{1/2} d\mu(t), \end{aligned}$$

$\alpha_0(t)$  is constant for  $t < -1$  and  $t > 1$ . If we choose  $\alpha_0(t)$  so that  $\alpha_0(1) = \alpha_0(1+0)$  and  $\alpha_0(-1) = \alpha_0(-1-0)$  then we can write

$$\int_{(-1,1)} U_{j+1}(t) (1-t^2)^{1/2} d\mu(t) = \int_{-1}^1 t^j d\alpha_0(t) \quad (8)$$

for all  $j \geq 0$ .

We show next that  $\mu$  is absolutely continuous on  $(-1, 1)$ . By (8) and the identity  $U_{j+3}(t) - U_{j+1}(t) = 2T_{j+3}(t)$  we have

$$2 \int_{(-1,1)} T_{j+3}(t) (1-t^2)^{1/2} d\mu(t) = -\int_{-1}^1 t^j (1-t^2) d\alpha_0(t) \quad (9)$$

for all  $j \geq 0$ . Hence for any  $n \geq 0$ ,

$$\begin{aligned} & \sum_{j=0}^n \left| \int_{(-1,1)} T_{j+3}(t)(1-t^2)^{1/2} d\mu(t) \right|^2 \\ & \leq \int_{-1}^1 \int_{-1}^1 \sum_{j=0}^n |st|^j (1-s^2)(1-t^2) d\alpha_0(s) d\alpha_0(t) \\ & \leq \left( \int_{-1}^1 d\alpha_0(t) \right)^2. \end{aligned}$$

Here we have used the inequality  $(1-uv)^{-1}(1-u^2)(1-v^2) < 1$ ,  $u, v \in (0, 1)$ .  
But

$$(1/\pi)^{1/2} T_0(t)(1-t^2)^{-1/4}, \quad (2/\pi)^{1/2} T_j(t)(1-t^2)^{-1/4}, \quad j = 1, 2, 3, \dots,$$

is a complete orthonormal system in  $L^2(-1, 1)$ , and so there exists an  $h \in L^2(-1, 1)$  such that

$$\int_{(-1,1)} T_j(t)(1-t^2)^{1/2} d\mu(t) = \int_{-1}^1 T_j(t)(1-t^2)^{-1/4} h(t) dt$$

for all  $j \geq 0$ . Then

$$\int_{(-1,1)} p(t)(1-t^2)^{1/2} d\mu(t) = \int_{-1}^1 p(t)(1-t^2)^{-1/4} h(t) dt$$

for every polynomial  $p(t)$ . It follows that, at least on  $(-1, 1)$ ,  $d\mu = f dt$  where  $f$  is real valued and integrable.

We assert that

$$f(x) = \lim_{r \uparrow 1} \frac{2}{\pi} \sum_{j=0}^{\infty} r^j U_j(x) \int_{-1}^1 U_j(t)(1-t^2)^{1/2} f(t) dt$$

a.e. on  $(-1, 1)$ . This follows from the fact that the Fourier sine series for the integrable function  $f(\cos \theta) \sin \theta$ ,  $0 < \theta < \pi$ , is Abel summable a.e. to the function [13, p. 99]. Since

$$\frac{1}{1-2xt+t^2} = \sum_{j=0}^{\infty} t^j U_j(x) \quad (10)$$

for  $-1 < t < 1$ ,  $-1 < x < 1$ , we obtain by (8)

$$\begin{aligned} f(x) &= p_0 + \lim_{r \uparrow 1} \frac{2r}{\pi} \int_{-1}^1 \sum_{j=0}^{\infty} (rt)^j U_{j+1}(x) d\alpha_0(t) \\ &= p_0 + \lim_{r \uparrow 1} \frac{2r}{\pi} \int_{-1}^1 \frac{2x - rt}{1 - 2rxt + r^2t^2} d\alpha_0(t) \\ &= p_0 + \frac{2}{\pi} \int_{-1}^1 \frac{2x - t}{1 - 2xt + t^2} d\alpha_0(t) \end{aligned}$$

a.e. on  $(-1, 1)$ . By redefining  $f$  on a set of measure zero we can assume that this representation is valid everywhere on  $(-1, 1)$ . Since

$$\frac{2x - t}{1 - 2xt + t^2} + \frac{t}{1 + t^2} = 2x \left(1 - \frac{2t}{1 + t^2} x\right)^{-1} (1 + t^2)^{-2}$$

we may rewrite the representation in the form

$$f(x) = p_1 + \int_{-1}^1 \frac{x}{1 - xt} d\alpha_1(t), \quad -1 < x < 1,$$

where  $p_1$  is a real constant and  $\alpha_1(t)$  is a nondecreasing function on  $[-1, 1]$ .

Now let  $a$  be an arbitrary positive number, and let  $b$  satisfy  $0 < b < a$ . By making a change of scale and using what was just proved we can argue that  $\mu$  is absolutely continuous on  $(-b, b)$ , and on  $(-b, b)$   $d\mu = f dx$  where

$$f(x) = p_b + \int_{-1/b}^{1/b} \frac{x}{1 - xt} d\alpha_b(t), \quad -b < x < b,$$

for some real constant  $p_b$  and nondecreasing function  $\alpha_b(t)$  on  $[-1/b, 1/b]$ . Hence  $\mu$  is (locally) absolutely continuous on  $(-a, a)$ . Since

$$p_b = f(0), \quad \alpha_b(1/b) - \alpha_b(-1/b) = f'(0)$$

a compactness argument shows that  $d\mu = f dx$  with  $f$  of the required form.

*Proof of sufficiency.* If  $\mu_1, \mu_2 \in \mathcal{U}$  and  $p_1, p_2$  are nonnegative numbers, then  $p_1\mu_1 + p_2\mu_2 \in \mathcal{U}$ . If  $(\mu_n)_1^\infty$  is a sequence in  $\mathcal{U}$  and if  $\mu$  is a real valued measure on  $(-a, a)$  such that

$$\lim_{n \rightarrow \infty} \int_{(-a, a)} \gamma d\mu_n = \int_{-a}^a \gamma d\mu$$

for all continuous functions  $\gamma$  which vanish off some closed subinterval of  $(-a, a)$ , then  $\mu \in \mathcal{U}$ . Therefore, it is enough to show that  $\mu \in \mathcal{U}$  if  $d\mu = f dx$

where  $f(x) = \pm 1$  or  $f(x) = x/(1 - \lambda x)$  for some  $\lambda$ ,  $-1/a < \lambda < 1/a$ . This follows from Proposition 1.

As dual problems we consider the following.

**PROBLEM 2.** *Determine the class  $\mathcal{V}_+$  of nonnegative lfs measures  $\nu$  on  $(-a, a)$  such that*

$$\int_{(-a,a)} |\tilde{\gamma}|^2 d\nu \leq \int_{(-a,a)} |\gamma|^2 d\nu \quad (11)$$

for all  $\gamma \in \mathcal{D}_a$ .

A similar problem is considered by Helson and Szegő [4].

**PROBLEM 3.** *Determine the class  $\mathcal{V}^-$  of lfs measures  $\nu$  on  $(-a, a)$  such that*

$$\begin{aligned} & \int_{(-b,b)} |\tilde{\gamma}(t)|^2 (a^2 - t^2)^{1/2} (b^2 - t^2)^{-1/2} d\nu(t) \\ & \leq \int_{(-b,b)} |\gamma(t)|^2 (a^2 - t^2)^{1/2} (b^2 - t^2)^{-1/2} d\nu(t) \end{aligned} \quad (12)$$

for all  $\gamma \in \mathcal{D}_a$  and numbers  $b$ ,  $0 < b < a$ , such that  $\gamma$  vanishes off  $(-b, b)$ .

It is not necessary to make any assumption regarding the existence of the integrals in (12). Our understanding of (12) is that the inequality holds for points  $b$  such that

$$\int_{(-b,b)} (b^2 - t^2)^{-1/2} d|\nu|(t) < \infty. \quad (13)$$

The excluded  $b$ 's form a Lebesgue null set. This follows from Fubini's theorem, since for any  $\epsilon$  and  $c$ ,  $0 < \epsilon < c < a$ , we have

$$\begin{aligned} & \int_{\epsilon}^c \left[ \int_{\epsilon < |t| < x} (x^2 - t^2)^{-1/2} d|\nu|(t) \right] dx \\ & = \int_{\epsilon < |t| < c} \left[ \int_{|t| < x < c} (x^2 - t^2)^{-1/2} dx \right] d|\nu|(t) \\ & = \int_{\epsilon < |t| < c} \log \frac{c + (c^2 - t^2)^{1/2}}{t} d|\nu|(t) < \infty. \end{aligned}$$

With this convention it is a consequence that for  $\nu \in \mathcal{V}^-$ , (13) holds for all  $b$ ,  $0 < b < a$ . See Theorem 2.

**PROPOSITION 2.** *Let  $\lambda$  be an lfs measure on  $(-a, a)$ . Suppose that for some fixed number  $b$ ,  $0 < b < a$ ,*

$$\int_{(-b,b)} |\tilde{\gamma}|^2 d\lambda \leq \int_{(-b,b)} |\gamma|^2 d\lambda \quad (14)$$



for all  $\gamma$  in  $\mathcal{D}_a$  which vanish off  $(-b, b)$ . Then on  $(-b, b)$ ,  $d\lambda = k dx$  where

$$k(x) = (b^2 - x^2)^{-1/2} \left[ p + qx - \int_{-1/b}^{1/b} \frac{x^2}{1 - xt} d\eta(t) \right], \quad (15)$$

$-b < x < b$ , for some real constants  $p = p_b$ ,  $q = q_b$  and nondecreasing function  $\eta(t) = \eta_b(t)$  on  $[-1/b, 1/b]$ .

*Proof.* The argument follows the proof of necessity of Theorem 1, the main change being that the roles of the polynomials  $T_n(x)$ ,  $U_n(x)$  are interchanged.

By making a change of scale we can assume that  $a > 1$  and  $b = 1$ . Substitute (7) into (14) to get

$$\sum_{j,k=0}^n \int_{(-1,1)} [U_j(t) U_k(t)(1 - t^2) - T_{j+1}(t) T_{k+1}(t)] d\lambda(t) c_j c_k^* \geq 0$$

and hence

$$-\sum_{j,k=0}^n \int_{(-1,1)} T_{j+k+2}(t) d\lambda(t) c_j c_k^* \geq 0$$

for any complex numbers  $c_0, \dots, c_n$ . As in the proof of Theorem 1 this implies that

$$-\int_{(-1,1)} T_{j+2}(t) d\lambda(t) = \int_{-1}^1 t^j d\eta_0(t) \quad (16)$$

for all  $j \geq 0$  where  $\eta_0(t)$  is a nondecreasing function on  $[-1, 1]$ . Since  $T_{j+2}(t) - T_j(t) = -2(1 - t^2) U_j(t)$  we obtain

$$-2 \int_{(-1,1)} U_{j+2}(t)(1 - t^2) d\lambda(t) = \int_{-1}^1 t^j (1 - t^2) d\eta_0(t)$$

analogous to (9), and from this we argue that  $\lambda$  is absolutely continuous on  $(-1, 1)$ . Let  $d\lambda = k dx$  on  $(-1, 1)$ .

From the Fourier cosine series for  $k(\cos \theta) \sin \theta$ ,  $0 < \theta < \pi$ , we obtain

$$(1 - x^2)^{1/2} k(x) = \frac{1}{\pi} \int_{-1}^1 T_0(t) k(t) dt + \lim_{r \uparrow 1} \frac{2}{\pi} \sum_{j=1}^{\infty} r^j T_j(x) \int_{-1}^1 T_j(t) k(t) dt$$

a.e. on  $(-1, 1)$ . Also, in place of (10) we have

$$\frac{1 - xt}{1 - 2xt + t^2} = \sum_{j=0}^{\infty} t^j T_j(x) \quad (17)$$

for  $-1 < t < 1$ ,  $-1 < x < 1$ . Now from (16) we obtain

$$\begin{aligned}(1-x^2)^{1/2} k(x) &= p_0 + q_0 x - \lim_{r \uparrow 1} \frac{2}{\pi} \sum_{j=0}^{\infty} r^{j+2} T_{j+2}(x) \int_{-1}^1 t' d\eta_0(t) \\ &= p_0 + q_0 x - \lim_{r \uparrow 1} \frac{2r^2}{\pi} \int_{-1}^1 \frac{2x^2 - 1 - rtx}{1 - 2rtx + r^2 t^2} d\eta_0(t) \\ &= p_0 + q_0 x - \frac{2}{\pi} \int_{-1}^1 \frac{2x^2 - 1 - tx}{1 - 2tx + t^2} d\eta_0(t)\end{aligned}$$

where  $p_0, q_0$  are real constants. By redefining  $k(x)$  on a set of measure zero we may assume that this representation holds at every point  $x$  of  $(-1, 1)$ . The identity

$$\frac{2x^2 - 1 - tx}{1 - 2tx + t^2} = -\frac{1}{1+t^2} - \frac{3t+t^3}{(1+t^2)^2} x + \frac{2x^2}{1-2tx+t^2} \frac{1-t^2}{(1+t^2)^2}$$

is used to rewrite the representation in the form (15).

LEMMA 2. Let  $g$  be a real valued function in  $L^p(-b, b)$  where  $p > 1$ . Let  $dv = g dx$ ,  $d\mu = -\tilde{g} dx$  on  $(-b, b)$ . Then

$$\int_{(-b,b)} |\tilde{\gamma}|^2 dv \leq \int_{(-b,b)} |\gamma|^2 dv \quad (18)$$

for all  $\gamma \in \mathcal{D}_b$  if and only if

$$\operatorname{Re} \int_{(-b,b)} \tilde{\gamma} \gamma^* d\mu \leq 0 \quad (19)$$

for all  $\gamma \in \mathcal{D}_b$ . In this case, (18) and (19) hold for any  $\gamma \in \mathcal{D}_a$ ,  $a > b$ , such that  $\gamma$  vanishes off  $(-b, b)$ .

Proof. If  $\gamma \in \mathcal{D}_b$  then by (2) and (3),

$$\begin{aligned}2 \operatorname{Re} \int_{(-b,b)} \tilde{\gamma} \gamma^* d\mu &= - \int_{-b}^b (\tilde{\gamma} \gamma^* + \gamma \tilde{\gamma}^*) \tilde{g} dt \\ &= \int_{-b}^b (\tilde{\gamma} \gamma^* + \gamma \tilde{\gamma}^*)^{\sim} g dt \\ &= \int_{(-b,b)} (|\tilde{\gamma}|^2 - |\gamma|^2) dv.\end{aligned}$$

The first assertion is now clear. The second follows by approximation.

PROPOSITION 3. *We have*

$$\begin{aligned}\int_{-b}^b |\tilde{\gamma}(t)|^2 (b^2 - t^2)^{-1/2} dt &= \int_{-b}^b |\gamma(t)|^2 (b^2 - t^2)^{-1/2} dt \\ \int_{-b}^b |\tilde{\gamma}(t)|^2 t(b^2 - t^2)^{-1/2} dt &= \int_{-b}^b |\gamma(t)|^2 t(b^2 - t^2)^{-1/2} dt\end{aligned}$$

for any  $\gamma \in \mathcal{D}_b$ , and indeed for any  $\gamma \in \mathcal{D}_a$ ,  $a > b$ , such that  $\gamma$  vanishes off  $(-b, b)$ .

*Proof.* The conjugate of  $g(x) = (b^2 - x^2)^{-1/2}$ ,  $x \in (-b, b)$ , is 0 in  $(-b, b)$  by [3, p. 247]. The conjugate of  $g(x) = x(b^2 - x^2)^{-1/2}$ ,  $x \in (-b, b)$ , is 1 in  $(-b, b)$ . In either case (19) holds for  $d\mu = \pm \tilde{g} dx$  and any  $\gamma \in \mathcal{D}_b$ . By Lemma 2, (18) holds for  $d\nu = \pm g dx$  and any  $\gamma \in \mathcal{D}_a$ ,  $a > b$ , such that  $\gamma$  vanishes off  $(-b, b)$ . The result follows.

LEMMA 3. *If  $\lambda \in [-1/a, 1/a]$  and  $0 < b < a$ , then*

$$\begin{aligned}-\int_{-b}^b |\tilde{\gamma}(t)|^2 t^2(b^2 - t^2)^{-1/2} (1 - \lambda t)^{-1} dt \\ \leq -\int_{-b}^b |\gamma(t)|^2 t^2(b^2 - t^2)^{-1/2} (1 - \lambda t)^{-1} dt\end{aligned}$$

for any  $\gamma \in \mathcal{D}_a$  such that  $\gamma$  vanishes off  $(-b, b)$ .

*Proof.* We compute the conjugate of  $g(x) = -x^2(b^2 - x^2)^{-1/2}(1 - \lambda x)^{-1}$ ,  $x \in (-b, b)$ . By [3, p. 247]

$$PV \frac{1}{\pi} \int_{-b}^b \frac{1}{t - x} (b^2 - t^2)^{-1/2} dt = \begin{cases} 0, & |x| < b \\ -\operatorname{sgn} x(x^2 - b^2)^{-1/2}, & |x| > b \end{cases} \quad (20)$$

Hence if  $-b < x < b$ ,

$$\begin{aligned}PV \frac{1}{\pi} \int_{-b}^b \frac{1}{t - x} (b^2 - t^2)^{-1/2} (1 - \lambda t)^{-1} dt \\ = (1 - \lambda x)^{-1} PV \frac{1}{\pi} \int_{-b}^b \left[ \frac{1}{t - x} - \frac{1}{t - 1/\lambda} \right] (b^2 - t^2)^{-1/2} dt \\ = (1 - \lambda x)^{-1} \operatorname{sgn} \lambda (\lambda^{-2} - b^2)^{-1/2} \\ = \lambda(1 - b^2 \lambda^2)^{-1/2} / (1 - \lambda x)\end{aligned}$$

at least for  $\lambda \neq 0$ . By (20) the same formula holds for  $\lambda = 0$  as well. If  $\lambda \neq 0$  then

$$g(x) = (\lambda^{-2} + \lambda^{-1}x)(b^2 - x^2)^{-1/2} - \lambda^{-2}(b^2 - x^2)^{-1/2}/(1 - \lambda x),$$

$x \in (-b, b)$ , and hence

$$-\tilde{g}(x) = -\lambda^{-1} + \lambda^{-1}(1 - b^2\lambda^2)^{-1/2}/(1 - \lambda x),$$

$x \in (-b, b)$ . By Proposition 1, (19) holds for  $d\mu = -\tilde{g} dx$  and any  $\gamma \in \mathcal{D}_b$ . It holds when  $\lambda = 0$  by a separate argument. The result now follows from Lemma 2.

**THEOREM 2.** *An lfs measure  $\nu$  on  $(-a, a)$  belongs to the class  $\mathcal{V}$  if and only if  $d\nu = g dx$  where*

$$g(z) = (a^2 - z^2)^{-1/2} \left[ p + qz - \int_{-1/a}^{1/a} \frac{z^2}{1 - zt} d\beta(t) \right], \quad (21)$$

$z \notin (-\infty, -a] \cup [a, \infty)$ , for some real constants  $p, q$  and nondecreasing function  $\beta(t)$  on  $[-1/a, 1/a]$ .

*Proof.* Suppose  $\nu \in \mathcal{V}$ . Let  $b$  be any number in  $(0, a)$  such that (13) holds. Let  $\lambda$  be any real valued measure on  $(-a, a)$  such that on  $(-b, b)$ ,  $d\lambda = (a^2 - t^2)^{1/2} (b^2 - t^2)^{-1/2} d\nu$ . By Proposition 2,  $d\lambda = k dx$  on  $(-b, b)$  where  $k(x)$  has the form (15). Since the set of points  $b$  in  $(0, a)$  such that (13) holds has  $a$  as a limit point, it follows that  $d\nu = g dx$  on  $(-a, a)$ . Further, for points  $b$  arbitrarily near  $a$ ,

$$g(x) = (a^2 - x^2)^{-1/2} \left[ p_b + q_b x - \int_{-1/b}^{1/b} \frac{x^2}{1 - xt} d\beta_b(t) \right], \quad -b < x < b,$$

where  $p_b, q_b$  are real constants and  $\beta_b(t)$  is a non-decreasing function on  $[-1/b, 1/b]$ . A straightforward compactness argument shows that  $g$  can be written in the form (21).

The sufficiency follows from Proposition 3 and Lemma 3.

**PROPOSITION 4.** *An lfs measure  $\nu$  on  $(-a, a)$  belongs to the class  $\mathcal{V}_+$  if and only if  $d\nu = g dx$  where  $g$  is a nonnegative function on  $(-a, a)$  of the form*

$$g(x) = (a^2 - x^2)^{-1/2} \left[ p + qx - \int_{-1/a}^{1/a} \frac{x^2}{1 - xt} d\beta(t) \right] \quad (22)$$

where  $p, q$  are real constants and  $\beta(t)$  is a nondecreasing function on  $[-1/a, 1/a]$ . Thus  $\mathcal{V}_+$  coincides with the class of nonnegative measures in  $\mathcal{V}$ .

*Proof.* Suppose  $\nu \in \mathcal{V}_+$ . Then  $\nu$  satisfies the hypotheses of Proposition 2 for every  $b \in (0, a)$ . Hence  $d\nu = g dx$ , and for each fixed  $b$  there is a representation

$$g(x) = (b^2 - x^2)^{-1/2} \left[ p_b + q_b x - \int_{-1/b}^{1/b} \frac{x^2}{1 - xt} d\beta_b(t) \right],$$

$-b < x < b$ , where  $p_b, q_b$  are real constants and  $\beta_b(t)$  is a nondecreasing function on  $[-1/b, 1/b]$ . By a compactness argument  $g$  has a representation in the form (22). Since  $\nu$  is nonnegative, so is  $g$ .

Conversely let  $d\nu = g dx$  where  $g$  is a nonnegative function of the form (22). By Theorem 2,  $\nu \in \mathcal{V}$  and hence (12) holds for all  $\gamma \in \mathcal{D}_a$  and all  $b \in (0, a)$  such that  $\gamma$  vanishes off  $(-b, b)$ . Since  $\nu$  is nonnegative we may let  $b \uparrow a$  and apply Fatou's lemma to get (11). That is,  $\nu \in \mathcal{V}_+$ .

*Note added in proof.* In connection with the Loewner problem, see also W. F. Donoghue, Jr., "Monotone Matrix Functions and Analytic Continuation," Springer Verlag, New York-Heidelberg-Berlin, 1974.

## REFERENCES

1. J. BENDAT AND S. SHERMAN, Monotone and convex operator functions, *Trans. Amer. Math. Soc.* **79** (1955), 58-71.
2. A. ERDÉLYI ET AL., "Higher Transcendental Functions," Bateman Manuscript Project, Vol. 2, McGraw-Hill, New York, 1953.
3. A. ERDÉLYI ET AL., "Tables of Integral Transforms," Bateman Manuscript Project, Vol. 2, McGraw-Hill, New York, 1954.
4. H. HELSON AND G. SZEGÖ, A problem in prediction theory, *Ann. Mat. Pura Appl.* **51** (1960), 107-138.
5. A. KORÁNYI, Note on the theory of monotone operator functions, *Acta Sci. Math. (Szeged)* **16** (1955), 241-245.
6. A. KORÁNYI, On a theorem of Löwner and its connections with resolvents of selfadjoint transformations, *Acta Sci. Math. (Szeged)* **17** (1956), 63-70.
7. K. LÖWNER, Über monotone Matrixfunktionen, *Math. Z.* **38** (1934), 177-216.
8. B. SZ.-NAGY, Remarks to the preceding paper of A. Koranyi, *Acta Sci. Math. (Szeged)* **17** (1956), 71-75.
9. E. C. TITCHMARSH, "Theory of Fourier Integrals," Oxford University Press, New York, 1959.
10. F. G. TRICOMI, "Integral Equations," Interscience, New York, 1957.
11. D. V. WIDDER, "The Laplace Transform," Princeton University Press, Princeton, NJ, 1972.
12. E. P. WIGNER AND J. V. NEUMANN, Significance of Loewner's theorem in the quantum theory of collisions, *Ann. of Math.* **59** (1954), 418-433.
13. A. ZYGMUND, "Trigonometric Series," Second Ed. Cambridge University Press, New York, 1968.